




<b>Cat. No:</b>	AB-84172
<b>Size:</b>	100ug
<b>Clone:</b>	POLY
<b>Concentration:</b>	1mg/ml
<b>Host:</b>	Rb
<b>Isotype:</b>	IgG
<b>Immunogen:</b>	Synthesized peptide derived from human Acrp30.
<b>Reactivity:</b>	Hu, Ms, Rt
<b>Applications:</b>	Western Blot: 1:500-1:2000 ELISA: 1:10000
<b>Molecular Weight:</b>	30kDa
<b>Purification:</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Background:</b>	Specificity: Adipoq Polyclonal Antibody detects endogenous levels of Adipoq protein. Tissue Specificity: Synthesized exclusively by adipocytes and secreted into plasma.
<b>Form:</b>	Liquid
<b>Buffer:</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide
<b>Storage:</b>	Store at -20°. Avoid freeze / thaw cycles.



Western blot analysis of lysates from 1) HeLa, 2) VEC, 3) 293 cells, [Green] primary antibody was diluted at 1:1000, 4° overnight, secondary antibody was diluted at 1:10000, 37° 1 hour. [Red] Tubulin β Monoclonal Antibody(5G3) antibody was diluted at 1:5000 as loading control, 4° overnight, secondary antibody was diluted at 1:10000, 37° 1 hour.



Western Blot analysis of specific cells using Adipoq polyclonal antibody



Western Blot (WB) analysis of HeLa cells using Adipoq polyclonal antibody.

**For Research use only  
IMMUNOLOGICAL SCIENCES**