



Cat. No:	AB-83897
Size:	100ug
Clone:	POLY
Concentration:	1mg/ml
Host:	Rb
Isotype:	IgG
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 1-108 of human ATP5J
Reactivity:	Hu, Ms,Rt
Applications:	Western Blot: 1:200 - 1:1000 Immunofluorescence: 1:50 - 1:100
Molecular Weight:	13kDa
Purification:	Aff.Pur.
Synonyms:	ATP5J; ATP5; ATP5A; ATPM; CF6; F6; ATP synthase-coupling factor 6, mitochondrial
Background:	Mitochondrial ATP synthase catalyzes ATP synthesis, utilizing an electrochemical gradient of protons across the inner membrane during oxidative phosphorylation. It is composed of two linked multi-subunit complexes: the soluble catalytic core, F1, and the membrane-spanning component, Fo, which comprises the proton channel. The F1 complex consists of 5 different subunits (alpha, beta, gamma, delta, and epsilon) assembled in a ratio of 3 alpha, 3 beta, and a single representative of the other 3. The Fo complex has nine subunits (a, b, c, d, e, f, g, F6 and 8). This gene encodes the F6 subunit of the Fo complex. The F6 subunit is required for F1 and Fo interactions. Alternatively spliced transcript variants encoding different isoforms have been identified for this gene. This gene has 1 or more pseudogenes.
Form:	Liquid
Buffer:	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Storage:	Store at -20°C. Avoid freeze / thaw cycles.



Western blot analysis of extracts of various cell lines, using ATP5J antibody at 1:1000 dilution.
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution.



Immunofluorescence analysis of U2OS cells using ATP5J antibody at dilution of 1:100. Blue: DAPI for nuclear staining.

Lysates/proteins: 25ug per lane.
Blocking buffer: 3% nonfat dry milk in
TBST.
Detection: ECL Basic Kit.
Exposure time: 90s.

**For Research use only
IMMUNOLOGICAL SCIENCES**