

| | |
|--------------------------|--|
| Cat. No: | AB-81400 |
| Conjugate: | Unconjugated |
| Size: | 100 ug |
| Clone: | POLY |
| Concentration: | 1mg/ml |
| Host: | Rb |
| Isotype: | IgG |
| Immunogen: | A synthetic peptide of human BCAT1 |
| Reactivity: | Hu ,Ms |
| Applications: | Western Blot: 1:500 - 1:2000 |
| Molecular Weight: | 43KDa |
| Purification: | Aff. Pur. |
| Synonyms: | BCAT1;BCATC;BCT1;ECA39;MECA39;PNAS121;PP18 |

Background: This gene encodes the cytosolic form of the enzyme branched-chain amino acid transaminase. This enzyme catalyzes the reversible transamination of branched-chain alpha-keto acids to branched-chain L-amino acids essential for cell growth. Two different clinical disorders have been attributed to a defect of branched-chain amino acid transamination: hypervalinemia and hyperleucine-isoleucinemia. As there is also a gene encoding a mitochondrial form of this enzyme, mutations in either gene may contribute to these disorders. Alternatively spliced transcript variants have been described.

| | |
|-----------------|---|
| Form: | Liquid |
| Buffer: | PBS with 0.02% sodium azide,50% glycerol,pH7.3. |
| Storage: | Store at -20°C. Avoid freeze / thaw cycles. |



Western blot analysis of extracts of various cell lines, using BCAT1 antibody at 1:1000 dilution.
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution.
Lysates/proteins: 25ug per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL West Pico Plus.
Exposure time: 1s.

**For Research use only
IMMUNOLOGICAL SCIENCES**