

| | |
|--------------------------|---|
| Cat. No: | AB-83773 |
| Conjugate: | Unconjugated |
| Size: | 100 ug |
| Clone: | Poly |
| Concentration: | 1mg/ml |
| Host: | Rabbit |
| Isotype: | IgG |
| Immunogen: | The antiserum was produced against synthesized peptide derived from human Caspase 3. AA range:126- 175 |
| Reactivity: | HU, Ms,Rt |
| Applications: | Western Blot: 1:500-2000 Immunohistochemistry (paraffin embedded tissues) : 1:50-300 Immunofluorescence: 1:50-300 |
| Molecular Weight: | 17kDA cleaved & 34kDA non cleaved |
| Purification: | The antibody was affinity- purified from rabbit antiserum by affinity- chromatography using epitope- specific immunogen. |
| Synonyms: | CASP3; CPP32; Caspase-3; CASP-3; Apopain; Cysteine protease CPP32; CPP-32; Protein Yama; SREBP cleavage activity 1; SCA- 1 |
| Background: | <p>caspase 3(CASP3) Homo sapiens This gene encodes a protein which is a member of the cysteine- aspartic acid protease (caspase) family. Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes which undergo proteolytic processing at conserved aspartic residues to produce two subunits, large and small, that dimerize to form the active enzyme. This protein cleaves and activates caspases 6, 7 and 9, and the protein itself is processed by caspases 8, 9 and 10. It is the predominant caspase involved in the cleavage of amyloid-beta 4A precursor protein, which is associated with neuronal death in Alzheimer's disease. Alternative splicing of this gene results in two transcript variants that encode the same protein.</p> |
| Form: | Liquid |
| Buffer: | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Storage: | Store at -20°C. Avoid repeated freeze-thaw cycles. |

**For Research use only
IMMUNOLOGICAL SCIENCES**