

Product Data Sheet: CD56 (N-CAM) Rabbit Polyclonal Antibody

Cat. No: AB-90239

Conjugate: Unconjugated

Size: 100 ug Clone: **POLY Concentration:** 1mg/ml Host: Rabbit Isotype: IgG

Recombinant fusion protein containing a sequence corresponding to amino acids Immunogen:

759-858 of human NCAM1 / CD56.

Hu, Ms, Rt Reactivity:

Western Blot: 1:500 - 1:1000

Immunohistochemistry(paraffin-embedded tissues): 1:50 - 1:200 **Applications:**

Immunofluorescence: 1:50 - 1:200 Immunocytochemistry: 1:50 - 1:200

Molecular Weight: 120-220kDa

Purification:

Aff. Pur. **Synonyms:** CD56; NCAM; MSK39; NCAM1 / CD56

> NCAM is a membrane-bound glycoprotein that plays a role in cell-cell and cellmatrix adhesion through both its homophilic and heterophilic binding activity. The neural cell adhesion molecule appears on early embryonic cells and is important in the formation of cell collectives and their boundaries at sites of morphogenesis. Later in development it is found on various differentiated tissues and is a major

Background: CAM mediating adhesion among neurons and between neurons and muscle.

NCAM gene is mapped to 11q23. The neural cell adhesion molecule (NCAM) can influence a number of diverse intercellular events, including junctional

communication, the association of axons with pathways and targets, and signals


that alter levels of neurotransmitter enzymes.

Form: Liquid


Buffer: PBS with 0.09% Sodium azide,50% glycerol,pH7.3.

At + 4°C for short term storage and at -20°C for longer time. Avoid Freeze and Storage:


Thaw cycles.



Western blot analysis of extracts of various cell lines, using NCAM1/CD56 antibody at 1:1000 dilution.



Immunohistochemistry of paraffinembedded human stomach using NCAM1 / CD56 antibody at dilution of 1:100



Immunofluorescence analysis of C6 cells using NCAM1 / CD56 antibody at dilution of 1:100. Blue: DAPI for nuclear staining.



Product Data Sheet: CD56 (N-CAM) Rabbit Polyclonal Antibody

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10,000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Pico Kit Exposure time: 90s

For Research use only IMMUNOLOGICAL SCIENCES