




| | |
|--------------------------|--|
| Cat. No: | AB-84755 |
| Conjugate: | Unconjugated |
| Size: | 100 ug |
| Clone: | POLY |
| Concentration: | 1mg/ml |
| Host: | Rabbit |
| Isotype: | IgG |
| Immunogen: | The antiserum was produced against synthesized peptide derived from the Internal region of human CD68. AA range:171-220. |
| Reactivity: | Human |
| Applications: | Western Blot: 1/500 - 1/2000. Immunohistochemistry (paraffin-embedded tissues) 1:100-300 ELISA: 1/20000. |
| Molecular Weight: | 37kD |
| Purification: | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Synonyms: | CD68; Macrosialin; Gp110; CD68 |
| Background: | This gene encodes a 110-kD transmembrane glycoprotein that is highly expressed by human monocytes and tissue macrophages. It is a member of the lysosomal/endosomal-associated membrane glycoprotein (LAMP) family. The protein primarily localizes to lysosomes and endosomes with a smaller fraction circulating to the cell surface. It is a type I integral membrane protein with a heavily glycosylated extracellular domain and binds to tissue- and organ-specific lectins or selectins. The protein is also a member of the scavenger receptor family. Scavenger receptors typically function to clear cellular debris, promote phagocytosis, and mediate the recruitment and activation of macrophages. Alternative splicing results in multiple transcripts encoding different isoforms. |
| Form: | Liquid |
| Buffer: | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Storage: | Store at -20°C. Avoid repeated freeze-thaw cycles. |



Western Blot analysis of mouse spleen cells using CD68 Polyclonal Antibody. Secondary antibody was diluted at




Immunohistochemical analysis of paraffin-embedded human-skin, antibody was diluted at 1:100.



Immunohistochemical analysis of paraffin-embedded human-skin, antibody was diluted at 1:100.

1:20000



Immunohistochemical analysis of
paraffin-embedded human-liver,
antibody was diluted at 1:100.

**For Research use only
IMMUNOLOGICAL SCIENCES**