

Cat. No:	MABN83864
Conjugate:	Unconjugated
Size:	100µL
Clone:	Monoclonal
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	A synthesized peptide derived from human CDC27
Reactivity:	Human,Mouse,Rat
Applications:	WB 1:1000-1:2000
Molecular Weight:	Calculated MW: 92 kDa ; Observed MW: 91 kDa
Purification:	Affinity Purification
Synonyms:	CDC27Hs; Cell division cycle protein 27 homolog; HNUC; APC3; ANAPC3;;CDC27
Background:	Component of the anaphase promoting complex/cyclosome (APC/C), a cell cycle-regulated E3 ubiquitin ligase that controls progression through mitosis and the G1 phase of the cell cycle. The APC/C complex acts by mediating ubiquitination and subsequent degradation of target proteins: it mainly mediates the formation of 'Lys-11'-linked polyubiquitin chains and, to a lower extent, the formation of 'Lys-48'- and 'Lys-63'-linked polyubiquitin chains.
Form:	Liquid
Buffer:	Purified antibody in PBS with 0.05% sodium azide,0.05% BSA and 50% glycerol.
Storage:	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.

**For Research use only
IMMUNOLOGICAL SCIENCES**