

<b>Cat. No:</b>	AB-84347
<b>Size:</b>	100ug
<b>Clone:</b>	POLY
<b>Concentration:</b>	1mg/ml
<b>Host:</b>	Rb
<b>Isotype:</b>	IgG
<b>Immunogen:</b>	The antiserum was produced against synthesized peptide derived from human PARP. AA range:196-245
<b>Reactivity:</b>	Hu, Ms
<b>Applications:</b>	WB 1:1000 Immunohistochemistry 1 :50-1:200
<b>Molecular Weight:</b>	89, 113kDa
<b>Purification:</b>	The antibody was affinity-purified from rabbit antiserum by affinity chromatography using epitope-specific immunogen.
<b>Background:</b>	<p>poly(ADP-ribose) polymerase 1(PARP1) Homo sapiens This gene encodes a chromatinassociated enzyme, poly(ADP-ribosyl)transferase, which modifies various nuclear proteins by poly(ADPribosyl) ation. The modification is dependent on DNA and is involved in the regulation of various important cellular processes such as differentiation, proliferation, and tumor transformation and also in the regulation of the molecular events involved in the recovery of cell from DNA damage. In addition, this enzyme may be the site of mutation in Fanconi anemia, and may participate in the pathophysiology of type I diabetes. The MW band of the cleaved form is recognized at 89kDa and the non-cleaved form is recognized at 113kDa.</p>
<b>Form:</b>	Liquid
<b>Buffer:</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Storage:</b>	20°C/1 year

**For Research use only  
IMMUNOLOGICAL SCIENCES**