

|                          |  |
|--------------------------|--|
| <b>Cat. No:</b>          | ABN09087   |
| <b>Conjugate:</b>        | Unconjugated   |
| <b>Size:</b>             | 100µL  |
| <b>Clone:</b>            | Polyclonal   |
| <b>Concentration:</b>    | 1mg/ml   |
| <b>Host:</b>             | Rabbit   |
| <b>Isotype:</b>          | IgG  |
| <b>Immunogen:</b>        | The antiserum was produced against synthesized peptide derived from human CKLF4. AA range:161-210  |
| <b>Reactivity:</b>       | Human,Mouse  |
| <b>Applications:</b>     | WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:100-1:300,ELISA 1:10000-1:20000   |
| <b>Molecular Weight:</b> | 26kDa  |
| <b>Purification:</b>     | Affinity purification  |
| <b>Synonyms:</b>         | CMTM4; CKLFSF4; CKLF-like MARVEL transmembrane domain-containing protein 4; Chemokine-like factor superfamily member 4   |
| <b>Background:</b>       | This gene belongs to the chemokine-like factor gene superfamily, a novel family that is similar to the chemokine and the transmembrane 4 superfamilies of signaling molecules. This gene is one of several chemokine-like factor genes located in a cluster on chromosome 16. Alternatively spliced transcript variants encoding different isoforms have been identified. [provided by RefSeq, Jul 2008],similarity:Belongs to the chemokine-like factor family.,similarity:Contains 1 MARVEL domain.,tissue specificity:Highly expressed in testis and prostate., |
| <b>Form:</b>             | Liquid   |
| <b>Buffer:</b>           | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.   |
| <b>Storage:</b>          | Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.   |

**For Research use only  
IMMUNOLOGICAL SCIENCES**