

<b>Cat. No:</b>	ABN09468
<b>Conjugate:</b>	Unconjugated
<b>Size:</b>	100µL
<b>Clone:</b>	Polyclonal
<b>Concentration:</b>	1mg/ml
<b>Host:</b>	Rabbit
<b>Isotype:</b>	IgG
<b>Immunogen:</b>	The antiserum was produced against synthesized peptide derived from human TAIP-2. AA range:171-220
<b>Reactivity:</b>	Human,Mouse,Monkey
<b>Applications:</b>	WB 1:500-1:2000,ELISA 1:10000-1:20000
<b>Molecular Weight:</b>	64kDa
<b>Purification:</b>	Affinity purification
<b>Synonyms:</b>	CSRNP3; FAM130A2; TAIP2; Cysteine/serine-rich nuclear protein 3; CSRNP-3; Protein FAM130A2; TGF-beta-induced apoptosis protein 2; TAIP-2
<b>Background:</b>	function: Binds to the consensus sequence 5'-AGAGTG-3' and has transcriptional activator activity. Plays a role in apoptosis., sequence caution: Unusual initiator. The initiator methionine is coded by a non-canonical CTG leucine codon., similarity: Belongs to the AXUD1 family., function: Binds to the consensus sequence 5'-AGAGTG-3' and has transcriptional activator activity. Plays a role in apoptosis., sequence caution: Unusual initiator. The initiator methionine is coded by a non-canonical CTG leucine codon., similarity: Belongs to the AXUD1 family.,
<b>Form:</b>	Liquid
<b>Buffer:</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
<b>Storage:</b>	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.

**For Research use only  
IMMUNOLOGICAL SCIENCES**