

| | |
|--------------------------|--|
| Cat. No: | MABN02961 |
| Conjugate: | Unconjugated |
| Size: | 100 μ L |
| Clone: | Monoclonal |
| Concentration: | 1mg/ml |
| Host: | Rabbit |
| Isotype: | IgG |
| Immunogen: | A synthetic peptide of human Cyclin H |
| Reactivity: | Human,Mouse |
| Applications: | WB 1:500-1:1000,ICC/IF 1:50-1:200,IP 1:20-1:50 |
| Molecular Weight: | Calculated MW: 38 kDa; Observed MW: 38 kDa |
| Purification: | Affinity Purification |
| Synonyms: | CCNH; Cyclin-H; MO15-associated protein; p34; p37 |
| Background: | Regulates CDK7, the catalytic subunit of the CDK-activating kinase (CAK) enzymatic complex. CAK activates the cyclin-associated kinases CDK1, CDK2, CDK4 and CDK6 by threonine phosphorylation. CAK complexed to the core-TFIID basal transcription factor activates RNA polymerase II by serine phosphorylation of the repetitive C-terminal domain (CTD) of its large subunit (POLR2A), allowing its escape from the promoter and elongation of the transcripts. Involved in cell cycle control and in RNA transcription by RNA polymerase II. Its expression and activity are constant throughout the cell cycle. |
| Form: | Liquid |
| Buffer: | 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA |
| Storage: | Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles. |

**For Research use only
IMMUNOLOGICAL SCIENCES**