

## Product Data Sheet: DDIT3

**Cat. No:** AB-83999

Conjugate: Unconjugated

Size: 100 ug

Clone: POLY

Concentration: 1mg/ml

Host: Rb

Isotype: IgG

**Immunogen:**Recombinant fusion protein containing a sequence corresponding to amino acids

1-288 of human HO-1

Reactivity: Hu, Ms, Rt

Western Blot: 1:500 - 1:2000

**Applications:** Immunohistochemistry: 1:50 – 1:200

Immunofluorescence: 1:50 - 1:200

**Molecular Weight:** 32kDa **Purification:** Aff. Pur.

**Synonyms:** HO-1;HMOX1D;HO-1;HSP32;bK286B10

Heme oxygenase, an essential enzyme in heme catabolism, cleaves heme to form biliverdin, which is subsequently converted to bilirubin by biliverdin

reductase, and carbon monoxide, a putative neurotransmitter. Heme oxygenase activity is induced by its substrate heme and by various nonheme substances.

Heme oxygenase occurs as 2 isozymes, an inducible heme oxygenase-1 and a constitutive heme oxygenase-2. HMOX1 and HMOX2 belong to the heme


oxygenase family.

Form: Liquid

**Background:** 


**Buffer:** PBS with 0.02% sodium azide,50% glycerol,pH7.3.

**Storage:** Store at -20°C. Avoid freeze / thaw cycles.




Western blot analysis of extracts of various cell lines, using HO-1 antibody at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution.

Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in



Immunohistochemistry of paraffin embedded human liver using HO-1 antibody at dilution of 1:200 (40x lens).




Immunofluorescence analysis of HeLa cells using HO-1 Polyclonal Antibody at dilution of 1:100 (40x lens).

Blue: DAPI for nuclear staining.



## Product Data Sheet: DDIT3

Detection: ECL West Pico Plus Exposure time: 10s.



Review for HO-1 Rabbit Polyclonal Antibody Experiment Type Western blot (WB)

## References

Product: HO-1 Rabbit pAb

Journal: British Journal Of Pharmacology

Application: WB IF:5.487 Species: Mus musculus

PMID: 26566260

Title: Trimetazidine prevents macrophage-mediated septic myocardial dysfunction via activation of the histone deacetylase

sirtuin 1

Product:HO-1 Rabbit pAb

Journal:Biochemical And Biophysical Research Communications

Application:WB IF:2.466 Species:Mus musculus

PMID:26903296

Title:Mst1 regulates hepatic lipid metabolism by inhibiting Sirt1 ubiquitination in mice References for HO-1 Rabbit pAb

Product: HO-1 Rabbit pAb

Journal:European Journal of Pharmacology Application:WB F:2.896 Species:Mus musculus

PMID:26973173

Title:Diabetes blocks the cardioprotective effects of sevoflurane postconditioning by impairing Nrf2/Brg1/HO-1 signaling

Product: HO-1 Rabbit pAb

Journal: Free Radical Biology and Medicine

Application: WB IF: 5.605 Species: Rattus norvegicus

PMID: 27840317

Title: Connexin43 regulates high glucose-induced expression of fibronectin, ICAM-1 and TGF-β1 via Nrf2/ARE pathway in

glomerular mesangial cells

Product: HO-1 Rabbit pAb Journal: Marine Drugs

Application: WB IF: 3.503 Species: Rattus norvegicus

PMID: 29621130

Title: Astaxanthin Promotes Nrf2/ARE Signaling to Inhibit HG-Induced Renal Fibrosis in GMCs

For Research use only IMMUNOLOGICAL SCIENCES