

Cat. No: AB-11173

Conjugate: Unconjugated

Size: 100 ul

Clone: POLY

Concentration: 1mg/ml

Host: Rb

Isotype: IgG

Reactivity: Hu, Rt, Ms

Applications: Elisa, Western Blot, Immunocytochemistry, Immunohistochemistry (frozen tissues)
Immunocytochemistry: 1:5000 dilution (PLP fixed rat brain sections) Western Blot: whole rat brain homogenate resulted in a pair of bands being detected at ~48 and ~51 kD at 1:1000-1:10000 with ECL Detection process. In some cases, it is required a higher dilution. For IHC (F) & ICC it is important to fix the cells or tissues with the PLP fixative (Please see protocol below)

Purification: Serum

Background: The antiserum was raised in a rabbit which was immunized with D2 (272-282) covalently attached onto a carrier protein, and it has been characterized by immunocytochemical, western immunoblot and ELISA techniques. The antiserum has been found to be highly specific for this peptide sequence and is suitable for immunocytochemical detection of both the short and the long forms of the D2 dopamine receptor. Rehydrate the antibody with 0.1 ml of PBS which contains 10 mg/ml BSA or with additional buffer (such as 10 mg/ml BSA in PBS) for more dilute antibody. 100% cross reactivity with D2 Dopamine Receptor (272-282), ~ 70% with D2S Dopamine Receptor, ~ 60% with D2L No cross reactivity with the other Dopamine Receptors.

Form: Liquid

**For Research use only
IMMUNOLOGICAL SCIENCES**