

| | |
|--------------------------|---|
| Cat. No: | ABN10293 |
| Conjugate: | Unconjugated |
| Size: | 100µL |
| Clone: | Polyclonal |
| Concentration: | 1mg/ml |
| Host: | Rabbit |
| Isotype: | IgG |
| Immunogen: | Synthesized peptide derived from part region of human protein |
| Reactivity: | Human,Mouse |
| Applications: | WB 1:500-1:2000,ELISA 1:5000-1:20000 |
| Molecular Weight: | 102kDa |
| Purification: | Affinity purification |

| | |
|--------------------|--|
| Background: | <p>Quality control in the endoplasmic reticulum (ER) ensures that only properly folded proteins are retained in the cell through recognition and degradation of misfolded or unassembled proteins. EDEM3 belongs to a group of proteins that accelerate degradation of misfolded glycoproteins in the ER (Hirao et al., 2006 [PubMed 16431915]).[supplied by OMIM, Mar 2008],domain:Contains a protease-associated domain (PA) of unknown function.,function:Involved in endoplasmic reticulum-associated degradation (ERAD). Accelerates the glycoprotein ERAD by proteasomes. This process depends on mannose-trimming from the N-glycans. Seems to have alpha 1,2-mannosidase activity.,PTM:N-glycosylated.,similarity:Belongs to the glycosyl hydrolase 47 family.,similarity:Contains 1 PA (protease associated) domain.,</p> |
| Form: | Liquid |
| Buffer: | Liquid in PBS containing 50% glycerol, and 0.02% New type preservative N. |
| Storage: | Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles. |

**For Research use only
IMMUNOLOGICAL SCIENCES**