

<b>Cat. No:</b>	AB-90101
<b>Conjugate:</b>	Unconjugated
<b>Size:</b>	100 ug
<b>Clone:</b>	POLY
<b>Concentration:</b>	1mg/ml
<b>Host:</b>	Rb
<b>Isotype:</b>	IgG
<b>Immunogen:</b>	Recombinant fusion protein containing a sequence corresponding to amino acids 143-288 of human FGF2
<b>Reactivity:</b>	Hu,Ms,Rt
<b>Applications:</b>	Western Blot : 1:500-1:2,000 Immunofluorescence/Immunocytochemistry: 1:50-1:200 Immunohistochemistry (paraffin, formalin, frozen Tissues): 1:50-1:200
<b>Molecular Weight:</b>	20kDa
<b>Purification:</b>	Aff. Pur.
<b>Background:</b>	<p>The protein encoded by this gene is a member of the fibroblast growth factor (FGF) family. FGF family members bind heparin and possess broad mitogenic and angiogenic activities. This protein has been implicated in diverse biological processes, such as limb and nervous system development, wound healing, and tumor growth. The mRNA for this gene contains multiple polyadenylation sites, and is alternatively translated from non-AUG (CUG) and AUG initiation codons, resulting in five different isoforms with distinct properties. The CUG-initiated isoforms are localized in the nucleus and are responsible for the intracrine effect, whereas, the AUG-initiated form is mostly cytosolic and is responsible for the paracrine and autocrine effects of this FGF.</p>
<b>Form:</b>	Liquid
<b>Buffer:</b>	Liquid in PBS with 0,02 % Sodium Azide and 50% glycerol pH 7.3.
<b>Storage:</b>	Storage for short term at 4°C recommended, for longer term at -20°C, minimize freeze/thaw cycles

**For Research use only  
IMMUNOLOGICAL SCIENCES**