

Cat. No:	MABN80772
Conjugate:	Unconjugated
Size:	100µL
Clone:	Monoclonal
Concentration:	1mg/ml
Host:	Mouse
Isotype:	Mouse IgG1
Immunogen:	Purified recombinant extracellular fragment of human FLT1 fused with hlgGfc tag expressed in HEK293 cells.
Reactivity:	Human,Mouse,Rat
Applications:	ELISA 1:5000-1:20000,FC 1:200-1:400
Molecular Weight:	100kDa
Purification:	Affinity Purification
Synonyms:	FLT; VEGFR1
Background:	<p>Fms-related tyrosine kinase 1 (vascular endothelial growth factor/vascular permeability factor receptor), also known as FLT1 or VEGFR-1. It is a member of the vascular endothelial growth factor receptor (VEGFR) family. VEGFR family members are receptor tyrosine kinases (RTKs) which contain an extracellular ligand-binding region with seven immunoglobulin (Ig)-like domains, a transmembrane segment, and a tyrosine kinase (TK) domain within the cytoplasmic domain. This protein binds to VEGFR-A, VEGFR-B and placental growth factor and plays an important role in angiogenesis and vasculogenesis. Expression of this receptor is found in vascular endothelial cells, placental trophoblast cells and peripheral blood monocytes. Multiple transcript variants encoding different isoforms have been found for this gene. Isoforms include a full-length transmembrane receptor isoform and shortened, soluble isoforms. The soluble isoforms are associated with the onset of pre-eclampsia.</p>
Form:	Liquid
Buffer:	Ascitic fluid containing 0.03% sodium azide.
Storage:	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.

**For Research use only
IMMUNOLOGICAL SCIENCES**