



<b>Cat. No:</b>	AB-84406
<b>Size:</b>	100ug
<b>Clone:</b>	POLY
<b>Concentration:</b>	1mg/ml
<b>Host:</b>	Rb
<b>Isotype:</b>	IgG
<b>Immunogen:</b>	Recombinant fusion protein containing a sequence corresponding to amino acids 22-138 of human GIPR.
<b>Reactivity:</b>	Hu, Ms, Rt
<b>Applications:</b>	vWestern Blot: 1:500 - 1:2000 Immunohistochemistry: 1:50 - 1:200 Immunofluorescence: 1:50 - 1:200 Flow cytometry: 1:50 - 1:200
<b>Molecular Weight:</b>	45kDa
<b>Purification:</b>	Aff. Pur.
<b>Synonyms:</b>	GIPR;PGQTL2
<b>Background:</b>	This gene encodes a G-protein coupled receptor for gastric inhibitory polypeptide (GIP), which was originally identified as an activity in gut extracts that inhibited gastric acid secretion and gastrin release, but subsequently was demonstrated to stimulate insulin release in the presence of elevated glucose. Mice lacking this gene exhibit higher blood glucose levels with impaired initial insulin response after oral glucose load. Defect in this gene thus may contribute to the pathogenesis of diabetes.
<b>Form:</b>	Liquid
<b>Buffer:</b>	PBS with 0.02% sodium azide,50% glycerol,pH7.3.
<b>Storage:</b>	Store at -20°C. Avoid freeze / thaw cycles.




Western blot analysis of extracts of various cell lines, using GIPR antibody at 1:1000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL West Pico Plus.



Immunohistochemistry of paraffin embedded rat brain using GIPR Rabbit pAb at dilution of 1:200 (40x lens).



Immunofluorescence analysis of 293 cells using GIPR polyclonal antibody (1:100). Secondary Antibody: Alexa Fluor 488 conjugated Goat anti Rabbit IgG

Exposure time: 90s.

**For Research use only  
IMMUNOLOGICAL SCIENCES**