

Cat. No:	MABN84748
Conjugate:	Unconjugated
Size:	100µL
Clone:	Monoclonal
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	A synthesized peptide derived from human HIF PH4
Reactivity:	Human
Applications:	WB 1:1000-1:2000,IHC 1:100-1:200,FC 1:20-1:100,IP 1:20-1:50
Molecular Weight:	Calculated MW: 57 kDa ; Observed MW: 47 kDa
Purification:	Affinity Purification
Synonyms:	EGLN4 ; HIFPH4; Hypoxia inducible factor prolyl 4 hydroxylase; P4H with transmembrane domain; P4htm; PH4; PHD4; Proline 4 hydroxylase; Prolyl hydroxylase domain containing 4;;HIF PH4
Background:	Catalyzes the post-translational formation of 4-hydroxyproline in hypoxia-inducible factor (HIF) alpha proteins. Hydroxylates HIF1A at 'Pro-402' and 'Pro-564'. May function as a cellular oxygen sensor and, under normoxic conditions, may target HIF through the hydroxylation for proteasomal degradation via the von Hippel-Lindau ubiquitination complex.
Form:	Liquid
Buffer:	Purified antibody in PBS with 0.05% sodium azide,0.05% BSA and 50% glycerol.
Storage:	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.

**For Research use only
IMMUNOLOGICAL SCIENCES**