

<b>Cat. No:</b>	AB-E8591
<b>Conjugate:</b>	unconjugated
<b>Size:</b>	100 ug
<b>Clone:</b>	Polyclonal
<b>Concentration:</b>	1mg/ml
<b>Host:</b>	Rabbit
<b>Isotype:</b>	IgG
<b>Immunogen:</b>	Synthesized peptide derived from the Internal region of human Histone H2A. 20-50
<b>Reactivity:</b>	Human, Mouse, Rat
<b>Applications:</b>	Western Blot 1:500- 1:2000 IHC(p) 1:100-1:300 Elisa: 1:20000
<b>Molecular Weight:</b>	15kd
<b>Purification:</b>	affinity purified
<b>Synonyms:</b>	HIST1H2AG; H2AFP; HIST1H2AI; H2AFC; HIST1H2AK; H2AFD; HIST1H2AL; H2AFI; HIST1H2AM; H2AFN; Histone H2A type 1; H2A.1; Histone H2A/p
<b>Background:</b>	<p>Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which ,approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H2A family.</p> <p>Transcripts from this gene lack polyA tails but instead contain a palindromic termination element.</p>
<b>Form:</b>	Liquid
<b>Buffer:</b>	PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide
<b>Storage:</b>	Store at -20°C. Avoid repeated freeze-thaw cycles.

**For Research use only  
IMMUNOLOGICAL SCIENCES**