

<b>Cat. No:</b>	AB-82662
<b>Conjugate:</b>	Unconjugated
<b>Size:</b>	100 ug
<b>Clone:</b>	POLY
<b>Concentration:</b>	1mg/ml
<b>Host:</b>	Rb
<b>Isotype:</b>	IgG
<b>Immunogen:</b>	Recombinant fusion protein containing a sequence corresponding to amino acids 20-330 of human CD11B
<b>Reactivity:</b>	Hu,Ms,Rt
<b>Applications:</b>	Western Blot: 1:500 -1:1000 Immunohistochemistry: 1:50 -1:100 Immunofluorescence: 1:50 -1:100
<b>Molecular Weight:</b>	180kDa
<b>Purification:</b>	Aff. Pur.
<b>Synonyms:</b>	ITGAM; CD11B; CR3A; MAC-1; MAC1A; MO1A; SLEB6; integrin alpha-M
<b>Background:</b>	<p>This gene encodes the integrin alpha M chain. Integrins are heterodimeric integral membrane proteins composed of an alpha chain and a beta chain. This I-domain containing alpha integrin combines with the beta 2 chain (ITGB2) to form a leukocytespecific integrin referred to as macrophage receptor 1 ('Mac-1'), or inactivated-C3b (iC3b) receptor 3 ('CR3'). The alpha M beta 2 integrin is important in the adherence of neutrophils and monocytes to stimulated endothelium, and also in the phagocytosis of complement coated particles. Multiple transcript variants encoding different isoforms have been found for this gene.</p>
<b>Form:</b>	Liquid
<b>Buffer:</b>	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
<b>Storage:</b>	Store at -20°C. Avoid freeze / thaw cycles.

**For Research use only  
IMMUNOLOGICAL SCIENCES**