

|                          |  |
|--------------------------|--|
| <b>Cat. No:</b>          | MAB-94656  |
| <b>Conjugate:</b>        | Unconjugated   |
| <b>Size:</b>             | 100 ug   |
| <b>Clone:</b>            | F10  |
| <b>Concentration:</b>    | 1mg/ml   |
| <b>Host:</b>             | Rb   |
| <b>Isotype:</b>          | IgG  |
| <b>Immunogen:</b>        | The antiserum was produced against synthesized peptide derived from the Internal region of human IL10. AA range:71-120   |
| <b>Reactivity:</b>       | Hu, Ms, Rt   |
| <b>Applications:</b>     | Western Blot: 1/500 - 1/2000 Immunohistochemistry (paraffin embedded tissues): 1:100-300 ELISA: 1/20000. IF 1:100-300  |
| <b>Molecular Weight:</b> | 20kDa  |
| <b>Purification:</b>     | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  |
| <b>Synonyms:</b>         | IL10; Interleukin-10; IL-10; Cytokine synthesis inhibitory factor; CSIF  |
| <b>Background:</b>       | <p>The protein encoded by this gene is a cytokine produced primarily by monocytes and to a lesser extent by lymphocytes. This cytokine has pleiotropic effects in immunoregulation and inflammation. It down-regulates the expression of Th1 cytokines, MHC class II Ags, and costimulatory molecules on macrophages. It also enhances B cell survival, proliferation, and antibody production. This cytokine can block NF-kappa B activity, and is involved in the regulation of the JAK-STAT signaling pathway. Knockout studies in mice suggested the function of this cytokine as an essential immunoregulator in the intestinal tract. Mutations in this gene are associated with an increased susceptibility to HIV-1 infection and rheumatoid arthritis</p> |
| <b>Form:</b>             | Liquid   |
| <b>Buffer:</b>           | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  |
| <b>Storage:</b>          | Store at -20°C. Avoid repeated freeze-thaw cycles.   |

**For Research use only  
IMMUNOLOGICAL SCIENCES**