




Cat. No:	MAB-94656
Conjugate:	Unconjugated
Size:	100 ug
Clone:	F10
Concentration:	1mg/ml
Host:	Rb
Isotype:	IgG
Immunogen:	The antiserum was produced against synthesized peptide derived from the Internal region of human IL10. AA range:71-120
Reactivity:	Hu, Ms, Rt
Applications:	Western Blot: 1/500 - 1/2000 Immunohistochemistry (paraffin embedded tissues): 1:100-300 ELISA: 1/20000. IF 1:100-300
Molecular Weight:	20kDa
Purification:	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Synonyms:	IL10; Interleukin-10; IL-10; Cytokine synthesis inhibitory factor; CSIF
Background:	The protein encoded by this gene is a cytokine produced primarily by monocytes and to a lesser extent by lymphocytes. This cytokine has pleiotropic effects in immunoregulation and inflammation. It down-regulates the expression of Th1 cytokines, MHC class II Ags, and costimulatory molecules on macrophages. It also enhances B cell survival, proliferation, and antibody production. This cytokine can block NF-kappa B activity, and is involved in the regulation of the JAK-STAT signaling pathway. Knockout studies in mice suggested the function of this cytokine as an essential immunoregulator in the intestinal tract. Mutations in this gene are associated with an increased susceptibility to HIV-1 infection and rheumatoid arthritis
Form:	Liquid
Buffer:	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Storage:	Store at -20°C. Avoid repeated freeze-thaw cycles.



Western Blot analysis of HepG2, Mouse liver cells using IL-10 Polyclonal Antibody. Secondary antibody was diluted at 1:20000




Immunofluorescence analysis of A549.
1, primary Antibody (red) was diluted at 1:200 (4°C overnight).
2, Goat Anti Rabbit IgG (H&L) - Alexa Fluor 594 Secondary antibody was diluted at 1:1000 (room temperature),



Immunohistochemical analysis of paraffin-embedded human breast cancer, antibody was diluted at 1:100

50min).3, Picture B: DAPI(blue) 10min.



Immunohistochemical analysis of
paraffin-embedded Human Cerebral
cortex.

- 1, Antibody was diluted at 1:200(4°
overnight).
- 2, High-pressure and temperature EDTA,
pH8.0 was used for antigen retrieval.
- 3,Secondary antibody was diluted at
1:200(room temperature, 30min).

**For Research use only
IMMUNOLOGICAL SCIENCES**