

Product Data Sheet: Interleukin-6

Cat. No: MAB-94635
Conjugate: Unconjugated

Size: 100 ug

Clone: 1621

Concentration: 1mg/ml

Host: Rabbit

Isotype: IgG

Immunogen: The antiserum was produced against synthesized peptide derived from the

Internal region of human IL6. AA range:131-180

Reactivity: Hu,Ms, Rt

Applications: Western Blot: 1/500 - 1/2000. Immunohistochemistry (paraffin-ebedded tissues):

1:100-300 Immunofluorescence: 1:100-300 ELISA: 1/20000

Molecular Weight: 23kDa

Purification: The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Synonyms: IL6; IFNB2; Interleukin-6; IL-6; B-cell stimulatory factor 2; BSF-2; CTL

differentiation factor; CDF; Hybridoma growth factor; Interferon beta-2; IFN-beta-2

This gene encodes a cytokine that functions in inflammation and the maturation of B cells. In addition, the encoded protein has been shown to be an endogenous pyrogen capable of inducing fever in people with autoimmune diseases or infections. The protein is primarily produced at sites of acute and chronic

Background: inflammation, where it is secreted into the serum and induces a transcriptional

inflammatory response through interleukin 6 receptor, alpha. The functioning of this gene is implicated in a wide variety of inflammation-associated disease states, including suspectibility to diabetes mellitus and systemic juvenile rheumatoid arthritis. Alternative splicing results in multiple transcript variants.

Form: Liquid

Buffer: Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Storage: Store at -20°C. Avoid freeze / thaw cycles.