


<b>Cat. No:</b>	MAB-94638
<b>Size:</b>	100 ug
<b>Clone:</b>	81E11
<b>Concentration:</b>	1mg/ml
<b>Host:</b>	Rb
<b>Isotype:</b>	IgG
<b>Immunogen:</b>	A synthesized peptide derived from human JNK
<b>Reactivity:</b>	Hu, Ms, Rt
<b>Applications:</b>	Western Blot: 1:1000 Immunofluorescence: 1:50 - 1:200
<b>Molecular Weight:</b>	46, 54 kDa


**Purification:** Monoclonal Antibodies are produced by immunizing animals with a recombinant human JNK2 fusion protein. Antibodies are purified by protein A and peptide affinity chromatography.

**Background:** The stress-activated protein kinase/Jun-aminoterminal kinase SAPK/JNK is potently and preferentially activated by a variety of environmental stresses including UV and gamma radiation, ceramides, inflammatory cytokines, and in some instances, growth factors and GPCR agonists (1-6). As with the other MAPKs, the core signaling unit is composed of a MAPKKK, typically MEKK1-MEKK4, or by one of the mixed lineage kinases (MLKs), which phosphorylate and activate MKK4/7. Upon activation, MKKs phosphorylate and activate the SAPK/JNK kinase (2). Stress signals are delivered to this cascade by small GTPases of the Rho family (Rac, Rho, cdc42) (3). Both Rac1 and cdc42 mediate the stimulation of MEKKs and MLKs (3). Alternatively, MKK4/7 can be activated in a GTPase-independent mechanism via stimulation of a germinai center kinase (GCK} family member (4). There are three SAPK/JNK genes each of which undergoes alternative splicing, resulting in numerous isoforms (3). SAPK/JNK, when active as a dimer, can translocate to the nucleus and regulate transcription through its effects on c-Jun, ATF-2, and other transcription factors (3,5). SAPK/JNK Antibody detects endogenous levels of total JNK1, JNK2 or JNK3 protein.

<b>Form:</b>	Liquid
<b>Buffer:</b>	PBS with 0.02% sodium azide,0.05% BSA,50% glycerol,pH7.3.
<b>Storage:</b>	Store at -20. Avoid freeze / thaw cycles.



Western blot analysis of extracts of various cell lines,



Immunofluorescence analysis of mouse brain using JNK1/2/3 Rabbit mAb at

using JNK1/2/3 Rabbit mAb at 1:1000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution.

Lysates/proteins: 25ug per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL West Pico Plus.

Exposure time: 30s.

dilution of 1:100 (40x lens).  
Blue: DAPI for nuclear staining.