

Cat. No:	AB- 83936
Conjugate:	Unconjugated
Size:	100 ug
Clone:	POLY
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	Laminin 111 from mouse EHS sarcoma cells obtained from Sigma-Aldrich.
Reactivity:	Human, Rat, Mouse
Applications:	Western Blot: 1:5,000 Immunofluorescence: 1:1,000-5,000 Immunocytochemistry: 1:1,000-5,000 Immunohistochemistry: 1:1,000-5,000
Molecular Weight:	440kDa, 220kDa, 158kDa
Purification:	Aff. Pur.

Background:

Laminins are high-molecular weight proteins and major components of the extracellular matrix. They are an important part of the basal lamina a protein network typically separating cells and tissues of different embryonic origin. Laminin was first isolated biochemically from Engelbreth-Holm-Swarm (EHS) mouse sarcoma cells, a cell line which produces large amounts of extracellular matrix material (1,2). Laminins are heterotrimeric proteins that contain one of five α -chains, one of three β -chains and one of three γ -chains (3-4). The EHS sarcoma cell derived laminin was originally named laminin 1, composed of $\alpha 1\beta 1\gamma 1$ polypeptides (5), though a later nomenclature named this form laminin 111, the numbers indicating the content of α , β and γ gene products (6). The distribution of the different laminin isoforms is developmental time and tissue-specific and laminin-111 expressed in the embryonic epithelium, but is also expressed in adult kidney, liver, testis, ovaries and brain blood vessels (7). Genetic ablation of laminin-111 results in embryonic death (8), and point mutations in the human $\alpha 1$ gene resulted in Poretti Boltshauser Syndrome, a rare disorder associated with cerebellar abnormalities, ataxia, cognitive problems and other issues. Injection of laminin-111 into muscle appeared to be a promising therapy in a mouse model Duchenne muscular dystrophy (DMD, 9). However transgenic overexpression of the laminin $\alpha 1$ protein appeared to have no beneficial affect also in a mouse DMD model (10). The Laminin antibody was made against laminin 111 isolated biochemically from mouse Engelbreth-Holm-Swarm (EHS) sarcoma cells (1,2). The antibody recognizes the several different laminin polypeptides on western blots and is a very useful reagent for visualizing basal lamina in mature and developing tissues. It is particularly useful in the CNS, allowing visualization of the basal lamina of endothelia, an important part of the blood brain barrier. Mouse select image above left for larger view.

Form:	Liquid
Buffer:	Affinity purified antibody at 1mg/mL in 50% PBS, 50% glycerol plus 5mM Na ₃
Storage:	Storage for short term at 4°C recommended, for longer term at -20°C.

**For Research use only
IMMUNOLOGICAL SCIENCES**