

Cat. No: AB-83938
Conjugate: Unconjugated
Size: 100 ug
Clone: POLY
Concentration: 1mg/ml
Host: Rb
Isotype: IgG
Immunogen: Recombinant human projection domain sequences, amino acids 377-1505.
Reactivity: Hu, Ms, Rt

Applications: Western Blot: 1:1,000
Immunofluorescence: 1:500-1:1000
Immunocytochemistry: 1;500-1:1000
Immunohistochemistry: 1:500-1:1000

Molecular Weight: ~280kDa

Purification: Purified


Background:

Microtubules are 25nm diameter protein rods found in most kinds of eukaryotic cells and are associated with a family of proteins called microtubule associated proteins (MAPs). MAPs play a crucial role in the regulation of microtubule dynamics and interactions in vivo. MAP2 was originally named as one of the higher molecular weight MAPs with an SDS-PAGE molecular weight of about 280kDa (1-3). There is a single mammalian MAP2 gene which may generate two high molecular weight proteins of ~280kDa name MAP2A and MAP2B and multiple lower molecular weight forms usually named MAP2C and MAP2D which run on SDS-PAGE gels at 60-70kDa. The lower molecular weight forms are found in neurons early in development, but later are replaced by the higher molecular weight forms (2). The MAP2A and MAP2B forms include a protein insert missing in MAP2C and MAP2D which forms fine filamentous protrusions from the sides of brain microtubules referred to as the "projection domain". This antibody was made against the projection domain sequences and so is specific for MAP2A and MAPB. MAP2 isoforms are expressed only in neuronal perikarya and dendrites so appropriate antibodies are used to identify neurons and dendrites in cell culture and sections. The MAP2 antibody was made against a mixture of recombinant human projection domain sequences, amino acids 377-1505. It binds to the MAP2A and MAP2B isoforms but not the MAP2C and MA2PD which lack projection domain sequences.


Form: Liquid

Buffer: Purified antibody at 1mg/mL in 50% PBS, 50% glycerol plus 5mM Na₃

Storage: Stable at 4°C for one year, for longer term store at -20°C



Western blot analysis of whole brain lysates using goat pAb to MAP2, MAP2, dilution 1:2,000 in green: [1] protein standard (red), [2] rat, and [3] mouse brain lysate. A band at about 280kDa corresponds to the MAP2A and MAP2B proteins.



Immunofluorescent analysis of rat brain stem section stained with goat pAb to MAP2, dilution 1:1,000 in red, and costained with mouse mAb to MBP, 7D2, dilution 1:2,500, in green. Following transcardial perfusion of rat with 4% paraformaldehyde, brain was post fixed for 24 hours, cut to 45 μ M, and free-floating sections were stained with above antibodies. The MAP2 antibody labels MAP2 protein in the perikarya and dendrites of the most neurons, notably motorneurons in the brain stem, and the MBP antibody stains myelin sheath around axons.

**For Research use only
IMMUNOLOGICAL SCIENCES**