

Cat. No:	AB-80184
Size:	100µg
Clone:	POLY
Concentration:	1mg/ml
Host:	Rb
Isotype:	IgG
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 24-99 of human MCP-1
Reactivity:	Hu, Ms
Applications:	Western Blot: 1:500 - 1:2000 Immunohistochemistry 1:100 - 1:200 Immunofluorescence: 1:50 - 1:100
Molecular Weight:	11kDa
Purification:	Aff. Pur.
Synonyms:	CCL2;GDCF-2;HC11;HSMCR30;MCAF;MCP-1;MCP1;SCYA2 ;SMC-CF
Background:	This gene is one of several cytokine genes clustered on the q-arm of chromosome 17. Chemokines are a superfamily of secreted proteins involved in immunoregulatory and inflammatory processes. The superfamily is divided into four subfamilies based on the arrangement of N-terminal cysteine residues of the mature peptide. This chemokine is a member of the CC subfamily which is characterized by two adjacent cysteine residues. This cytokine displays chemotactic activity for monocytes and basophils but not for neutrophils or eosinophils. It has been implicated in the pathogenesis of diseases characterized by monocytic infiltrates, like psoriasis, rheumatoid arthritis and atherosclerosis. It binds to chemokine receptors CCR2 and CCR4
Form:	Liquid
Buffer:	PBS with 0.02% sodium azide,50% glycerol,pH7.3
Storage:	Store at -20°C. Avoid freeze / thaw cycles

**For Research use only
IMMUNOLOGICAL SCIENCES**