

Cat. No:	MABN85779
Conjugate:	Unconjugated
Size:	100µL
Clone:	Monoclonal
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	A synthetic peptide of human MEKK2
Reactivity:	Human,Mouse,Rat
Applications:	WB 1:500-1:1000,ICC 1:50-1:200,IP 1:10-1:20
Molecular Weight:	Calculated MW: 70 kDa; Observed MW: 70 kDa
Purification:	Affinity Purification
Synonyms:	MAP3K2; MAPKKK2; MEKK2; Mitogen-activated protein kinase kinase kinase 2; MAPK/ERK kinase kinase 2; MEK kinase 2; MEKK 2
Background:	<p>The protein encoded by this gene is a member of serine/threonine protein kinase family. This kinase preferentially activates the kinases involved in the MAP kinase signaling pathway including MAPK7 and MAP2K4. This kinase has been shown to directly phosphorylate and activate I kappa B kinases (IKKs), and thus plays a role in NF-kappa B signaling pathway. This kinase has also been found to bind and activate protein kinase C-related kinase 2 (PRKCL2/PRK2), which suggests its involvement in PRKCL2 regulated signaling process.</p>
Form:	Liquid
Buffer:	Purified antibody in TBS with 0.05% sodium azide,0.05%BSA and 50% glycerol.
Storage:	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.

**For Research use only
IMMUNOLOGICAL SCIENCES**