

Product Data Sheet: MET (c-Met)

Isotype: IgG

Immunogen:Recombinant fusion protein containing a sequence corresponding to amino acids

25-325 of human MET.

Reactivity: Hu,Ms,Rt

Applications: Western Blot: 1:500 -1:2000

Molecular Weight: 155-180kDa

Purification: Aff. Pur.

Synonyms: MET; AUTS9; DFNB97; HGFR; RCCP2; c-Met; hepato cyte growth factor receptor

This gene encodes a member of the receptor tyrosine kinase family of proteins and the product of the proto-oncogene MET. The encoded preproprotein is proteolytically processed to generate alpha and beta subunits that are linked via disulfide bonds to form the mature receptor. Further processing of the beta subunit results in the formation of the M10 peptide, which has been shown to reduce lying fibracia. Pinding of its ligand, beneficiate growth factor, induces

Background: reduce lung fibrosis. Binding of its ligand, hepatocyte growth factor, induces dimerization and activation of the receptor, which plays a role in cellular survival,

embryogenesis, and cellular migration and invasion. Mutations in this gene are associated with papillary renal cell carcinoma, hepatocellular carcinoma, and various head and neck cancers. Amplification and overexpression of this gene are

also associated with multiple human cancers.

Form: Liquid

Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Storage: Store at -20°C. Avoid freeze / thaw cycles.