






Cat. No:	MAB-94594
Conjugate:	Unconjugated
Size:	100ug
Clone:	ARC0879
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	A synthetic peptide corresponding to a sequence within amino acids 1-100 of human Myelin oligodendrocyte glycoprotein
Reactivity:	Hu, Ms, Rt
Applications:	Western Blot: 1:500 – 1:1000 Immunofluorescence: 1:50 – 1:200 Immunocytochemistry: 1:50 – 1:200
Molecular Weight:	28kDa
Purification:	Affinity purification
Synonyms:	BTN6; BTNL11; MOGIG2; NRCLP7
Background:	The product of this gene is a membrane protein expressed on the oligodendrocyte cell surface and the outermost surface of myelin sheaths. Due to this localization, it is a primary target antigen involved in immune-mediated demyelination. This protein may be involved in completion and maintenance of the myelin sheath and in cell-cell communication. Alternatively spliced transcript variants encoding different isoforms have been identified.
Form:	Liquid
Buffer:	PBS with 0.02% sodium azide, 0.05% BSA, 50% glycerol, pH 7.3.
Storage:	Store at -20°C. Avoid freeze / thaw cycles.



Western blot analysis of extracts of various cell lines, using Myelin oligodendrocyte glycoprotein antibody at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL West Pico Plus. Exposure time: 10s.



Immunofluorescence analysis of rat brain using Myelin oligodendrocyte glycoprotein antibody at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.



Immunofluorescence analysis of mouse brain using Myelin oligodendrocyte glycoprotein antibody at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.

**For Research use only
IMMUNOLOGICAL SCIENCES**