

Cat. No:	AB-84338
Conjugate:	Unconjugated
Size:	100 ug
Clone:	POLY
Concentration:	1mg/ml
Host:	Rb
Isotype:	IgG
Immunogen:	Nanoghomeobox
Reactivity:	Hu,Ms
Applications:	Western Blot: 1:500-1:2000 Immunofluorescence: 1:20-1:200
Molecular Weight:	43-48 kDa
Purification:	Aff. Pur. ≥95% as determined by SDS-PAGE

Background: Transcription regulator involved in inner cell mass and embryonic stem(ES) cells proliferation and self-renewal. Imposes pluripotency on ES cells and prevents their differentiation towards extraembryonic endoderm and trophectoderm lineages. Blocks bone morphogenetic protein-induced mesoderm differentiation of ES cells by physically interacting with SMAD1 and interfering with the recruitment of coactivators to the active SMAD transcriptional complexes. Acts as a transcriptional activator or repressor. Binds optimally to the DNA consensus sequence 5'-TAAT[GT][GT]-3' or 5'-[CG][GA][CG]C[GC]ATTAN[GC]-3'. Able to autorepress its expression in differentiating(ES) cells: binds to its own promoter following interaction with ZNF281/ZFP281, leading to recruitment of the NuRD complex and subsequent repression of expression. When overexpressed, promotes cells to enter into S phase and proliferation.

Form:	Liquid
Buffer:	PBS with 0.02% sodium azide and 50% glycerol pH 7.3
Storage:	T -20°C for 12 months (Avoid repeated freeze / thaw cycles.)

**For Research use only
IMMUNOLOGICAL SCIENCES**