

Cat. No:	MAB-94715
Conjugate:	Unconjugated
Size:	100 ug
Clone:	D15E13
Concentration:	1mg/ml
Host:	Mouse
Isotype:	IgG1
Immunogen:	Recombinant Protein
Reactivity:	Human, Mouse, Rat
Applications:	WesternBlot:1/500-2000 Immunoprecipitation:1/200 Immunofluorescence:1/100-200 Immunohistochemistry:1/200-500
Molecular Weight:	65kDa
Synonyms:	NFKB3 p65 RELA Transcription factor p65
Background:	NFkB p65 is ubiquitinated leading to its proteosomal degradation which is required for termination of the NFkB response. Phosphorylation of NFkB p65 on S536 stimulates acetylation of K310 by CBP enhancing transcriptional activity. NFkB p65 is also acetylated at K122 enhancing DNA binding and impairing the interaction with NFKBIA. The protein is deacetylated by HDAC3. Invasion of a host by a pathogen is frequently associated with the activation of NF-kB which coordinates various aspects of immune function required for resistance to infection.
Form:	Liquid
Buffer:	PBS with 0.02% sodium azide and 50% glycerol pH 7.4.
Storage:	Store at -20°C. Avoid repeated freeze-thaw cycles.

**For Research use only
IMMUNOLOGICAL SCIENCES**