

Cat. No:	ABN14658
Conjugate:	Unconjugated
Size:	100μL
Clone:	Polyclonal
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	The antiserum was produced against synthesized peptide derived from human NFRKB. AA range:463-512
Reactivity:	Human,Mouse,Monkey
Applications:	WB 1:500-1:2000,IHC 1:50-1:300,ELISA 1:2000-1:20000
Molecular Weight:	110kDa
Purification:	Affinity purification
Synonyms:	NFRKB; INO80G; Nuclear factor related to kappa-B-binding protein; DNA-binding protein R kappa-B; INO80 complex subunit G function: Binds to the DNA consensus sequence 5'-GGGGAATCTCC-3'.PTM: Phosphorylated upon DNA damage, probably by ATM or ATR.,subunit: Component of the chromatin-remodeling INO80 complex, at least composed of ACTL6A, ACTR5, ACTR8, RVBL1, RVBL2, INO80, INO80B, INO80C, INO80D and INO80E.,tissue specificity: Expressed in thymus, brain, testes, spleen and liver.,
Background:	function: Binds to the DNA consensus sequence 5'-GGGGAATCTCC-3'.PTM: Phosphorylated upon DNA damage, probably by ATM or ATR.,subunit: Component of the chromatin-remodeling INO80 complex, at least composed of ACTL6A, ACTR5, ACTR8, RVBL1, RVBL2, INO80, INO80B, INO80C, INO80D and INO80E.,tissue specificity: Expressed in thymus, brain, testes, spleen and liver.,
Form:	Liquid
Buffer:	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Storage:	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.

**For Research use only
IMMUNOLOGICAL SCIENCES**