

Product Data Sheet: NOTCH 3 Rabbit Polyclonal Antibody

Cat. No: AB-10483

Conjugate: Unconjugated

Size: 100µg Clone: **POLY Concentration:** 1mg/ml Host: Rabbit Isotype: IqG

A synthetic peptide corresponding to a sequence within amino acids 2200-2321 of Immunogen:

human NOTCH3.

Reactivity: Human, Mouse, Rat

Western Blot: 1:1000 - 1:5000 **Applications:**

Immunohistochemistry(paraffin-embedded tissues): 1:50 - 1:200

Molecular Weight: 90kDa

Purification: Affinity purification

IMF2; LMNS; CASIL; CADASIL; CADASIL1; NOTCH3 **Synonyms:**

> This gene encodes the third discovered human homologue of the Drosophilia melanogaster type I membrane protein notch. In Drosophilia, notch interaction with its cell-bound ligands (delta, serrate) establishes an intercellular signalling pathway that plays a key role in neural development. Homologues of the notchligands have also been identified in human, but precise interactions between

these ligands and the human notch homologues remains to be determined. Mutations in NOTCH3 have been identified as the underlying cause of cerebral

autosomal dominant arteriopathy with subcortical infarcts and


leukoencephalopathy (CADASIL).

Form: Liquid

Background:


Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Store at -20°C. Avoid freeze / thaw cycles. Storage:




Western blot analysis of lysates from K-562 cells, using NOTCH3 Rabbit pAb at 1:2000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST.



Immunohistochemistry analysis of paraffin-embedded mouse spleen using NOTCH3 Rabbit pAb at dilution of 1:100 (40x

lens).Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded mouse pancreas using NOTCH3 Rabbit pAb at dilution of 1:100 (40x

lens).Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Product Data Sheet: NOTCH 3 Rabbit Polyclonal Antibody

Detection: ECL West Pico Plus. Exposure time: 10s.

For Research use only IMMUNOLOGICAL SCIENCES