

Cat. No:	ABN15084
Conjugate:	Unconjugated
Size:	100µL
Clone:	Polyclonal
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	Synthesized peptide derived from human protein . at AA range: 90-170
Reactivity:	Human,Mouse
Applications:	WB 1:500-1:2000,ELISA 1:5000-1:20000
Molecular Weight:	20kDa
Purification:	Affinity purification

Background: The protein encoded by this gene belongs to the ornithine decarboxylase antizyme family, which plays a role in cell growth and proliferation by regulating intracellular polyamines. Expression of antizymes requires +1 ribosomal frameshifting, which is enhanced by high levels of polyamines. Antizymes in turn bind to and inhibit ornithine decarboxylase (ODC), the key enzyme in polyamine biosynthesis; thus, completing the auto-regulatory circuit. This gene encodes antizyme 2, the second member of the antizyme family. Like antizyme 1, antizyme 2 has broad tissue distribution, inhibits ODC activity and polyamine uptake, and stimulates ODC degradation in vivo; however, it fails to promote ODC degradation in vitro. Antizyme 2 is expressed at lower levels than antizyme 1, but is evolutionary more conserved, suggesting it likely has an important biological role. Studies also show different subalternative products:A ribosomal frameshift occurs between the codons for Ser-32 and Asp-33. An autoregulatory mechanism enables modulation of frameshifting according to the cellular concentration of polyamines,function: Binds to, and destabilizes, ornithine decarboxylase. Does not accelerate ornithine decarboxylase degeneration.,similarity: Belongs to the ODC antizyme family.,

Form:	Liquid
Buffer:	Liquid in PBS containing 50% glycerol, and 0.02% New type preservative N.
Storage:	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.

**For Research use only
IMMUNOLOGICAL SCIENCES**