

Cat. No: AB-10220**Size:** 200 µg**Clone:** POLY**Concentration:** 1mg/ml**Host:** Rb**Isotype:** IgG**Reactivity:** Hu**Applications:** WB, IP, IHC(P)**Purification:** Aff. Pur.**Synonyms:** Cyclin-dependent kinase inhibitor 2A; multiple tumor suppressor 1, ARF; CDK4I; CDKN2; CMM2; INK4; INK4A;MLM; MTS-1; MTS1; P14; P14ARF; P16; P16-INK4A; P16INK4; P16INK4A**Background:** The progression through the cell cycle is regulated by cyclins and their cognate Cdk by promoting cell cycle transitions (1,2,3). This orderly progression can be inhibited by a family of proteins known as CDK inhibitors (CDIs) that bind to cyclin/Cdk complexes and halt cell division (3). p21 (also designated WAF1/Cip1) is one has been shown to inhibit the activity of each member of the cyclin/Cdk family and over expression of this protein inhibits the proliferation of mammalian cells (5). The expression of p21 is inducible by a wide range of stress stimuli and its transcription can be enhanced by p53 (6). Another member of the CDIs is p27 (also known as Kip1), which also sees up regulation in response to antimitogenic stimuli (7). The increased protein expression of p27 results in cellular arrest by binding to cyclin/Cdk complexes, like cyclin D1/Cdk4 (4,8). An additional CDI has been found to bind Cdk4 and Cdk6, p16 (INK4A), and when such complexes are formed, the progression of the cell cycle is halted (9). It has become increasingly clear that p16 is a very important tumor suppressor gene whose frequent loss occurs early in many human cancers. p16 is a major target in carcinogenesis that is rivaled in frequency only by p53 (10).**Form:** Liquid**Buffer:** Each vial contains 200 µg/ 1 ml of affinity purified rabbit IgG, p16 (C20) AB-10220, in 1 ml PBS containing 0.1 % sodium azide and 0.2% gelatin.**Storage:** Store this product at 4° C, do not freeze. The product is stable for one year from the date of shipment.

Immunoperoxidase staining of formalin-fixed, paraffin-embedded human tonsil showing expression in the mantle zone and the majority of cells in the germinal center negative - dilution1:50

References

1. Harper JW, Adami GR, Wei N, Keyomarsi K, Elledge SJ. 1993. The p21 Cdk-interacting protein Cip1 is a potent inhibitor of

- G1 cyclindependent kinases. Cell 75(4):805-816.
2. Gartel AL, Serfas MS, Tyner AL. 1996. p21 - -negative regulator of the cell cycle. Proc Soc Exp Biol Med 213(2):138-149.
 3. Moller MB. 2000. p27 in cell cycle control and cancer. Leuk Lymphoma 39(1-2):19-27.
 4. Sgambato A, Cittadini A, Faraglia B, Weinstein IB. 2000. Multiple functions of p27(Kip1) and its alterations in tumor cells: a review. J Cell Physiol 183(1):18-27.
 5. Xiong Y, Hannon GJ, Zhang H, Casso D, Kobayashi R, Beach D. 1993. p21 is a universal inhibitor of cyclin kinases. Nature 366(6456):634.
 6. Gorospe M, Wang X, Holbrook NJ. 1999. Functional role of p21 during the cellular response to stress. Gene Expr 7(4-6):377-385.
 7. Slingerland J, Pagano M. 2000. Regulation of the cdk inhibitor p27 and its deregulation in cancer. J Cell Physiol 183(1):10-17.
 8. Toyoshima H, Hunter T. 1994. p27, a novel inhibitor of G1 cyclin-Cdk protein kinase activity, is related to p21. Cell 78(1):67-74.
 9. Shapiro GI, Edwards CD, Rollins BJ. 2000. The physiology of p16(INK4A)-mediated G1 proliferative arrest. Cell Biochem Biophys 33(2):189-197.
 10. Liggett WH Jr, Sidransky D. 1998. Role of the p16 tumor suppressor gene in cancer. J Clin Oncol 16(3):1197-1206.

**For Research use only
IMMUNOLOGICAL SCIENCES**