

Product Data Sheet: P63

Cat. No: AB-81811

Conjugate: Unconjugated

Size: 100 ug
Clone: POLY
Concentration: 1mg/ml
Host: Rb

Isotype: IgG

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids

421-680 of human p63.

Reactivity: Hu, Ms, Rt

Applications: Western Blot: 1:500 - 1:2000 Immunohistochemistry: 1:50 - 1:200

Molecular

Weight: 75kDa

Purification: Aff. Pur.

Synonyms: AIS;B(p51A);B(p51B);EEC3;KET;LMS;NBP;OFC8;RHS;SHFM4;TP53CP;TP53L;

TP73L;p40;p51;p53CP;p63;p73H;p73L;TP63

This gene encodes a member of the p53 family of transcription factors. The functional domains of p53 family proteins include an N-terminal transactivation domain, a central DNA-binding domain and an oligomerization domain. Alternative splicing of this gene and the use of alternative promoters results in multiple transcript variants encoding different isoforms that vary in their functional properties. These isoforms function during skin development and maintenance,

properties. These isoforms function during skin development and maintenance, adult stem/progenitor cell regulation, heart development and premature aging.

Some isoforms have been found to protect the germline by eliminating oocytes or testicular germ cells that have suffered DNA damage. Mutations in this gene are

testicular germ cells that have suffered DNA damage. Mutations in this gene are associated with ectodermal dysplasia, and cleft lip/palate syndrome 3 (EEC3); splithand/foot malformation 4 (SHFM4); ankyloblepharon-ectodermal defects-cleft lip/palate; ADULT syndrome (acro-dermatoungual- lacrimal-tooth); limb-mammary

syndrome; Rap-Hodgkin syndrome (RHS); and orofacial cleft 8.

Form: Liquid

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Storage: Store at -20°C. Avoid freeze / thaw cycles.