

## **Product Data Sheet:** P63

Cat. No: AB-81811

Conjugate: Unconjugated

Size: 100 ug Clone: **POLY Concentration:** 1mg/ml

Host: Rb Isotype: IqG

Recombinant fusion protein containing a sequence corresponding to amino acids Immunogen:

421-680 of human p63.

Reactivity: Hu, Ms, Rt

Western Blot: 1:500 - 1:2000 **Applications:** 

Immunohistochemistry: 1:50 - 1:200

Molecular

75kDa Weight:

**Purification:** Aff. Pur.

AIS;B(p51A);B(p51B);EEC3;KET;LMS;NBP;OFC8;RHS;SHFM4;TP53CP;TP53L; **Synonyms:** 

TP73L;p40;p51;p53CP;p63;p73H;p73L;TP63

This gene encodes a member of the p53 family of transcription factors. The functional domains of p53 family proteins include an N-terminal transactivation domain, a central DNA-binding domain and an oligomerization domain. Alternative splicing of this gene and the use of alternative promoters results in multiple transcript variants encoding different isoforms that vary in their functional properties. These isoforms function during skin development and maintenance,

**Background:** adult stem/progenitor cell regulation, heart development and premature aging. Some isoforms have been found to protect the germline by eliminating oocytes or testicular germ cells that have suffered DNA damage. Mutations in this gene are

associated with ectodermal dysplasia, and cleft lip/palate syndrome 3 (EEC3); splithand/foot malformation 4 (SHFM4); ankyloblepharon-ectodermal defects-cleft lip/palate; ADULT syndrome (acro-dermatoungual- lacrimal-tooth); limb-mammary

syndrome; Rap-Hodgkin syndrome (RHS); and orofacial cleft 8.

Form: Liquid

**Buffer:** PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Storage: Store at -20°C. Avoid freeze / thaw cycles.

×

Western blot analysis of extracts of various cell lines, using p63 antibody at 1:1000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG

(H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in

TBST.

Detection: ECL West Pico Plus.Exposure

×

Immunohistochemistry of paraffin embedded human esophageal cancer using p63 antibody at dilution of 1:50 (40x lens).



## Product Data Sheet: P63

time: 3min.

## For Research use only IMMUNOLOGICAL SCIENCES