




<b>Cat. No:</b>	ABP10261
<b>Conjugate:</b>	Unconjugated
<b>Size:</b>	100 ug
<b>Clone:</b>	Poly
<b>Concentration:</b>	1mg/ml
<b>Host:</b>	Rb
<b>Isotype:</b>	IgG
<b>Immunogen:</b>	Synthesized peptide derived from human c-Myc around the phosphorylation site of S62.
<b>Reactivity:</b>	Hu, Ms, Rt
<b>Applications:</b>	Western Blot: 1:500-1:2000 Immunohistochemistry: 1:100-1:300 Immunofluorescence: 1:200-1:1000 ELISA: 1:40000
<b>Molecular Weight:</b>	50 kDa
<b>Purification:</b>	The antibody was affinity-purified from rabbit antiserum by affinity chromatography using epitope-specific immunogen.
<b>Background:</b>	Transcription factor that binds DNA in a non-specific manner, yet also specifically recognizes the core sequence 5'-CAC[GA]TG-3'. Activates the transcription of growth-related genes. Phospho-c-Myc (S62) Polyclonal Antibody detects endogenous levels of c- Myc protein only when phosphorylated at S62.
<b>Form:</b>	liquid
<b>Buffer:</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide
<b>Storage:</b>	Store at -20°C. Avoid freeze / thaw cycles.



Western Blot (WB) analysis of 293T, AD293, 22RV1 and HeLa cells, using Anti-Phospho-c-Myc (S62) polyclonal antibody.



Western Blot (WB) analysis of 293 cells using Anti-Phospho-c-Myc (S62) polyclonal antibody



Immunohistochemical analysis of paraffin-embedded Human breast cancer using Anti-Phospho-c-Myc (S62) polyclonal antibody. Antibody was diluted at 1:100 (4°C, overnight). High-pressure and temperature Tris-EDTA, pH8.0 was used for antibody retrieval. Negative control (right) obtained from antibody was pre-absorbed by immunogen peptide.

**For Research use only  
IMMUNOLOGICAL SCIENCES**