




<b>Cat. No:</b>	ABP-0199
<b>Conjugate:</b>	Unconjugated
<b>Size:</b>	100 ug
<b>Clone:</b>	Poly
<b>Concentration:</b>	1mg/ml
<b>Host:</b>	Rb
<b>Isotype:</b>	IgG
<b>Immunogen:</b>	Synthesized peptide derived from human EGFR around the phosphorylation site of T693.
<b>Reactivity:</b>	Hu, Ms, Rt
<b>Applications:</b>	Western Blot: 1:500-1:2000 Immunohistochemistry: 1:100-1:300 Immunofluorescence: 1:200-1:1000 ELISA: 1:40000
<b>Purification:</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using a epitope-specific immunogen.
<b>Background:</b>	Phospho-EGFR (T693) polyclonal antibody detects endogenous levels of EGFR protein only when phosphorylated at T693.
<b>Form:</b>	liquid
<b>Buffer:</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Storage:</b>	Store at -20°C, and avoid repeat freeze-thaw cycles.




Western blot analysis of lysates from A431 cells, using EGFR (Phospho-Thr693) Antibody. The lane on the right is blocked with the phospho peptide.



Western blot analysis of various cells using Phospho-EGFR (T693) Polyclonal Antibody



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma, using EGFR (Phospho-Thr693) Antibody. The picture on the right is blocked with the phospho peptide.



Immunofluorescence analysis of HUVEC cells, using EGFR (Phospho-Thr693) Antibody. The lane on the right is blocked with the phospho peptide.

**For Research use only  
IMMUNOLOGICAL SCIENCES**