


<b>Cat. No:</b>	ABP-0026
<b>Conjugate:</b>	Unconjugated
<b>Size:</b>	100 ug
<b>Clone:</b>	Poly
<b>Concentration:</b>	1mg/ml
<b>Host:</b>	Rb
<b>Isotype:</b>	IgG
<b>Reactivity:</b>	Hu
<b>Applications:</b>	WB: 1:1000
<b>Molecular Weight:</b>	175 kDa

**Purification:** Polyclonal antibodies are produced by immunizing animals with a synthetic phosphopeptide corresponding to residues surrounding Tyr992 of human EGF receptor. Antibodies are purified by protein A and peptide affinity chromatography.

**Background:** The epidermal growth factor (EGF) receptor is a 170 kDa transmembrane tyrosine kinase that belongs to the HER/ErbB protein family. Ligand binding results in receptor dimerization, autophosphorylation, activation of downstream signaling, internalization and lysosomal degradation (1,2). Phosphorylation of EGF receptor (EGFR) at Tyr845 in the kinase domain is implicated in stabilizing the activation loop, maintaining the active state enzyme and providing a binding surface for substrate proteins (3,4). c-Src is involved in phosphorylation of EGFR at Tyr845 (5). The SH2 domain of PLC $\beta$  binds at phospho-Tyr992, resulting in activation of PLC $\beta$ -mediated downstream signaling (6). Phosphorylation of EGFR at Tyr1045 creates a major docking site for c-Cbl, an adaptor protein that leads to receptor ubiquitination and degradation following EGFR activation (7,8). The GRB2 adaptor protein binds activated EGFR at phospho-Tyr1068 (9). A pair of phosphorylated EGFR residues (Tyr1148 and Tyr1173) provides a docking site for the Shc scaffold protein, with both sites involved in MAP kinase signaling activation (2). Phosphorylation of EGFR at specific serine and threonine residues attenuates EGFR kinase activity. EGFR carboxy-terminal residues Ser1046 and Ser1047 are phosphorylated by CaM kinase II; mutation of either of these serines results in upregulated EGFR tyrosine autophosphorylation (10). Phospho-EGF Receptor (Tyr992) Antibody detects endogenous EGF receptor only when phosphorylated at tyrosine 992. The antibody may cross-react with other activated EGF receptor family members (e.g. ErbB2).

<b>Form:</b>	liquid
<b>Buffer:</b>	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
<b>Storage:</b>	Store at -20°C. Avoid freeze / thaw cycles



Western blot analysis of extracts of A431 cell line, using Phospho-EGFR (Tyr992) antibody.

## References

- (1) Hackel, P.O. et al. (1999) *Curr. Opin. Cell Biol.* 11, 184-189. (2) Zwick, E. et al. (1999) *Trends Pharmacol. Sci.* 20, 408-412. (3) Cooper, J.A. and Howell, B. (1993) *Cell* 73, 1051-1054. (4) Hubbard, S.R. et al. (1994) *Nature* 372, 746-754. (5) Biscardi, J.S. et al. (1999) *J. Biol. Chem.* 274, 8335-8343. (6) Emllet, D.R. et al. (1997) *J. Biol. Chem.* 272, 4079-4086. (7) Levkowitz, G. et al. (1999) *Mol. Cell* 4, 1029-1040. (8) Ettenberg, S.A. et al. (1999) *Oncogene* 18, 1855-1866. (9) Rojas, M. et al. (1996) *J. Biol. Chem.* 271, 27456-27461. (10) Feinmesser, R.L. et al. (1999) *J. Biol. Chem.* 274, 16168-16173.

**For Research use only  
IMMUNOLOGICAL SCIENCES**