

Cat. No:	MAB-94213
Conjugate:	Unconjugated
Size:	100 ug
Clone:	174A9
Concentration:	1mg/ml
Host:	Rb
Isotype:	IgG
Reactivity:	Hu, Ms, Rt
Applications:	WB 1:1000
Molecular Weight:	15 kDa


Purification: Monoclonal antibody is produced by immunizing animals with a synthetic phosphopeptide corresponding to residues surrounding Ser65 of mouse 4E-BP1 protein.

Background: Translation repressor protein 4E-BP1 (also known as PHAS-1) inhibits cap-dependent translation by binding to the translation initiation factor eIF4E. Hyperphosphorylation of 4E-BP1 disrupts this interaction and results in activation of cap-dependent translation (1). Both the PI3 kinase/Akt pathway and FRAP/mTOR kinase regulate 4E-BP1 activity (2,3). Multiple 4E-BP1 residues are phosphorylated in vivo (4). While phosphorylation by FRAP/mTOR at Thr37 and Thr46 does not prevent the binding of 4E-BP1 to eIF4E, it is thought to prime 4E-BP1 for subsequent phosphorylation at Ser65 and Thr70 (5). Phospho-4E-BP1 (Ser65) (174A9) Rabbit mAb recognizes endogenous levels of 4EBP1 protein only when phosphorylated at Ser65.

Form: liquid

Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Storage: Store at -20°C. Avoid freeze / thaw cycles.



Western blot analysis on 293 cell lysates
using Phospho-4E-BP1 (Ser65) antibody

References

(1) Pause, A. et al. (1994) Nature 371, 762-7. (2) Brunn, G.J. et al. (1997) Science 277, 99-101. (3) Gingras, A.C. et al. (1998)

Genes Dev 12, 502-13. (4) Fadden, P. et al. (1997) J Biol Chem 272, 10240-7. (5) Gingras, A.C. et al. (1999) Genes Dev 13, 1422-37..

**For Research use only
IMMUNOLOGICAL SCIENCES**