

Cat. No: ABP-0031
Conjugate: Unconjugated
Size: 100 ug
Clone: Poly
Concentration: 1mg/ml
Host: Rb
Isotype: IgG
Reactivity: Hu, Ms, Rt
Applications: WB: 1:1000
Molecular Weight: 15-20 kDa

Purification: Polyclonal antibodies are produced by immunizing animals with a synthetic phosphopeptide corresponding to the sequence of rat 4E-BP1. Antibodies are purified by protein A and peptide affinity chromatography.

Background: Translation repressor protein 4E-BP1 (also known as PHAS-1) inhibits cap-dependent translation by binding to the translation initiation factor eIF4E. Hyperphosphorylation of 4E-BP1 disrupts this interaction and results in activation of cap-dependent translation (1). Both the PI3 kinase/Akt pathway and FRAP/mTOR kinase regulate 4E-BP1 activity (2,3). Multiple 4E-BP1 residues are phosphorylated in vivo (4). While phosphorylation by FRAP/mTOR at Thr37 and Thr46 does not prevent the binding of 4E-BP1 to eIF4E, it is thought to prime 4E-BP1 for subsequent phosphorylation at Ser65 and Thr70 (5). Phospho-4E-BP1 (Thr70) Antibody detects endogenous levels of 4E-BP1 only when phosphorylated at threonine 70. This antibody does not detect 4E-BP1 phosphorylated at other sites.

Form: liquid
Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Storage: Store at -20°C. Avoid freeze / thaw cycles.

**For Research use only
IMMUNOLOGICAL SCIENCES**