


Cat. No:	ABP-0359
Conjugate:	Unconjugated
Size:	100 ug
Clone:	Poly
Concentration:	1mg/ml
Host:	Rb
Isotype:	IgG
Immunogen:	A phospho specific peptide corresponding to residues surrounding S632 of human HDAC4
Reactivity:	Hu, Ms, Rt
Applications:	WB 1:500-1:2000
Molecular Weight:	140kDa
Purification:	Affinity purification
Background:	Histones play a critical role in transcriptional regulation, cell cycle progression, and developmental events. Histone acetylation/deacetylation alters chromosome structure and affects transcription factor access to DNA. The protein encoded by this gene belongs to class II of the histone deacetylase/acuc/apha family. It possesses histone deacetylase activity and represses transcription when tethered to a promoter. This protein does not bind DNA directly, but through transcription factors MEF2C and MEF2D. It seems to interact in a multiprotein complex with RbAp48 and HDAC3.
Form:	liquid
Buffer:	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Storage:	Store at -20oC or -80oC. Avoid freeze / thaw cycles.



Western blot analysis of extracts from 293 cells using Phospho-HDAC4-S632 antibody

**For Research use only
IMMUNOLOGICAL SCIENCES**