

<b>Cat. No:</b>	ABP-0359
<b>Conjugate:</b>	Unconjugated
<b>Size:</b>	100 ug
<b>Clone:</b>	Poly
<b>Concentration:</b>	1mg/ml
<b>Host:</b>	Rb
<b>Isotype:</b>	IgG
<b>Immunogen:</b>	A phospho specific peptide corresponding to residues surrounding S632 of human HDAC4
<b>Reactivity:</b>	Hu, Ms, Rt
<b>Applications:</b>	WB 1:500-1:2000
<b>Molecular Weight:</b>	140kDa
<b>Purification:</b>	Affinity purification
<b>Background:</b>	<p>Histones play a critical role in transcriptional regulation, cell cycle progression, and developmental events. Histone acetylation/deacetylation alters chromosome structure and affects transcription factor access to DNA. The protein encoded by this gene belongs to class II of the histone deacetylase/acuc/apha family. It possesses histone deacetylase activity and represses transcription when tethered to a promoter. This protein does not bind DNA directly, but through transcription factors MEF2C and MEF2D. It seems to interact in a multiprotein complex with RbAp48 and HDAC3.</p>
<b>Form:</b>	liquid
<b>Buffer:</b>	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
<b>Storage:</b>	Store at -20oC or -80oC. Avoid freeze / thaw cycles.

**For Research use only  
IMMUNOLOGICAL SCIENCES**