

Cat. No:	MAB-94464
Conjugate:	Unconjugated
Size:	100 ug
Clone:	A09
Concentration:	1mg/ml
Host:	Rb
Isotype:	IgG
Immunogen:	A synthetic phosphorylated peptide around S139 of human Histone H2AX
Reactivity:	Hu
Applications:	Western Blot: 1:1000 Immunohistochemistry: 1:50 Immunofluorescence: 1:50
Molecular Weight:	17kDa
Purification:	Affinity purification
Synonyms:	H2AFX; H2A.X; H2A/X; H2AX; histone H2AX

Background: Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene encodes a replication-independent histone that is a member of the histone H2A family, and generates two transcripts through the use of the conserved stemloop termination motif, and the polyA addition motif.

Form:	liquid
Buffer:	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Storage:	Store at -20°C. Avoid freeze / thaw cycles.

**For Research use only
IMMUNOLOGICAL SCIENCES**