



Cat. No:	ABP11206
Conjugate:	Unconjugated
Size:	100 ug
Clone:	Poly
Concentration:	1mg/ml
Host:	Rb
Isotype:	IgG
Immunogen:	A phospho specific peptide corresponding to residues surrounding Y1349 of human MET
Reactivity:	Hu
Applications:	Western Blot: 1:500 - 1:2000 Immunohistochemistry 1:50 - 1:200 Immunofluorescence: 1:50 - 1:200
Molecular Weight:	155.541 kDa
Purification:	Affinity purification

Background: This gene encodes a member of the receptor tyrosine kinase family of proteins and the product of the proto-oncogene MET. The encoded preproprotein is proteolytically processed to generate alpha and beta subunits that are linked via disulfide bonds to form the mature receptor. Further processing of the beta subunit results in the formation of the M10 peptide, which has been shown to reduce lung fibrosis. Binding of its ligand, hepatocyte growth factor, induces dimerization and activation of the receptor, which plays a role in cellular survival, embryogenesis, and cellular migration and invasion. Mutations in this gene are associated with papillary renal cell carcinoma, hepatocellular carcinoma, and various head and neck cancers. Amplification and overexpression of this gene are also associated with multiple

Form:	liquid
Buffer:	PBS with 0.02% sodium azide, 50% glycerol, pH7.3
Storage:	Store at -20°C, and avoid repeat freeze-thaw cycles



Western blot analysis of extracts of HeLa cells, using Phospho-MET-Y1349 antibody at 1:1000 dilution. HeLa cells were treated by HGF (40ng/ml) for 10 minutes after serum-starvation overnight. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane.



Immunofluorescence analysis of U2OS cells using Phospho-MET-Y1349 antibody
Blue: DAPI for nuclear staining.

Blocking buffer: 3% BSA. Detection: ECL
Basic Kit. Exposure time: 90s.

**For Research use only
IMMUNOLOGICAL SCIENCES**