

Product Data Sheet: Phospho-p70 S6k (S235/236)

Cat. No: ABP-0538

Conjugate: Unconjugated

Size: 100 ug
Clone: Poly
Concentration: 1mg/ml

Host: Rb Isotype: IgG

Immunogen: A synthetic phosphorylated peptide around S235 & S236 of human RPS6.

Reactivity: Hu

Applications: Western Blot: 1:500 – 1:2000

Molecular Weight: 32KDa

Synonyms: RPS6;S6;RPS6

Ribosomes, the organelles that catalyze protein synthesis, consist of a small 40S subunit and a large 60S subunit. Together these subunits are composed of 4 RNA species and approximately 80 structurally distinct proteins. This gene encodes a cytoplasmic ribosomal protein that is a component of the 40S subunit. The protein belongs to the S6E family of ribosomal proteins. It is the major substrate of protein kinases in the ribosome, with subsets of five C-terminal serine residues phosphorylated by different protein kinases. Phosphorylation is induced by a wide

Background: phosphorylated by different protein kinases. Phosphorylation is induced b range of stimuli, including growth factors, tumor-promoting agents, and


mitogens. Dephosphorylation occurs at growth arrest. The protein may contribute to the control of cell growth and proliferation through the selective translation of particular classes of mRNA. As is typical for genes encoding ribosomal proteins, there are multiple processed pseudogenes of this gene dispersed through the

genome.

Form: liquid

Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Storage: Store at -20°C. Avoid freeze / thaw cycles.



Western blot analysis of extracts of various cell lines, using Phospho-S6 Ribosomal Protein (RPS6)-S235/236 antibody at 1:1000 dilution.HeLa cells NIH/3T3 cells and PC-12 cells were treated by 10% FBS at 37°C for 30 minutes after serum-starvation overnight.NIH/3T3 cells were treated by



Product Data Sheet: Phospho-p70 S6k (S235/236)

PDGF (100 ng/ml) at 37°C for 30 minutes after serum-starvation overnight.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution.

Lysates/proteins: 25ug per lane.

Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL West Pico Plus.

Exposure time: 1s.

For Research use only IMMUNOLOGICAL SCIENCES