

<b>Cat. No:</b>	MAB-94331
<b>Conjugate:</b>	Unconjugated
<b>Size:</b>	100 ul
<b>Clone:</b>	D8C2Z
<b>Concentration:</b>	1mg/ml
<b>Host:</b>	Rb
<b>Isotype:</b>	IgG
<b>Reactivity:</b>	Hu, Ms, Rt, Monkey
<b>Applications:</b>	Western Blotting 1:1000 Immunohistochemistry 1:100-1:300 Immunoprecipitation: 1:200-500 ELISA: 1:10000
<b>Molecular Weight:</b>	86 kDa
<b>Purification:</b>	Monoclonal antibody is produced by immunizing animals with a synthetic phospho-peptide corresponding to residues surrounding Ser727 of human Stat3 protein.
<b>Background:</b>	The Stat3 transcription factor is an important signaling molecule for many cytokines and growth factor receptors (1) and is required for murine fetal development (2). Research studies have shown that Stat3 is constitutively activated in a number of human tumors (3,4) and possesses oncogenic potential (5) and anti-apoptotic activities (3). Stat3 is activated by phosphorylation at Tyr705, which induces dimerization, nuclear translocation, and DNA binding (6,7). Transcriptional activation seems to be regulated by phosphorylation at Ser727 through the MAPK or mTOR pathways (8,9). Stat3 isoform expression appears to reflect biological function as the relative expression levels of Stat3 $\alpha$ (86 kDa) and Stat3 $\beta$ (79 kDa) depend on cell type, ligand exposure, or cell maturation stage (10). It is notable that Stat3 $\beta$ lacks the serine phosphorylation site within the carboxy-terminal transcriptional activation domain (8). Phospho-Stat3 (Ser727) (D8C2Z) Rabbit mAb recognizes endogenous levels of Stat3 protein only when phosphorylated at Ser727
<b>Form:</b>	liquid
<b>Storage:</b>	Store at -20°C and avoid repeat freeze-thaw cycles.

**For Research use only**  
**IMMUNOLOGICAL SCIENCES**