


Cat. No: ABP-4002
Conjugate: Unconjugated
Size: 100 ug
Clone: Poly
Concentration: 1mg/ml
Host: Rb
Isotype: IgG
Reactivity: Hu, Ms, Rt, Monkey
Applications: Western Blotting 1:1000 Immunohistochemistry 1:100-1:300
 Immunoprecipitation: 1:200-500 ELISA: 1:10000
Molecular Weight: 86 kDa


Purification: Polyclonal antibody is produced by immunizing animals with a synthetic phosphopeptide corresponding to residues surrounding Ser727 of human Stat3 protein.

Background: The Stat3 transcription factor is an important signaling molecule for many cytokines and growth factor receptors (1) and is required for murine fetal development (2). Research studies have shown that Stat3 is constitutively activated in a number of human tumors (3,4) and possesses oncogenic potential (5) and anti-apoptotic activities (3). Stat3 is activated by phosphorylation at Tyr705, which induces dimerization, nuclear translocation, and DNA binding (6,7). Transcriptional activation seems to be regulated by phosphorylation at Ser727 through the MAPK or mTOR pathways (8,9). Stat3 isoform expression appears to reflect biological function as the relative expression levels of Stat3 β (86 kDa) and Stat3 α (79 kDa) depend on cell type, ligand exposure, or cell maturation stage (10). It is notable that Stat3 α lacks the serine phosphorylation site within the carboxy-terminal transcriptional activation domain (8). Phospho-Stat3 (Ser727) Rabbit pAb recognizes endogenous levels of Stat3 protein only when phosphorylated at Ser727

Form: liquid
Buffer: Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Storage: Store at -20°C, and avoid repeat freeze-thaw cycles.



Western blot analysis of extracts from serum-starved U266 cells, untreated (-) or treated with Human Interferon- γ (hIFN- γ ; 50ng/ml, 15 min; +), HeLa cells, untreated (-) or treated with Human Tumor Necrosis Factor- α (hTNF- α ;



Western blot analysis of extracts from serum-starved A-431 cells, untreated (-) or treated with Human Epidermal Growth Factor(hEGF; 100ng/ml, 30 min; +) and SH-PPase (SH-PPase; +) using Phospho-Stat3 (Ser727) Rabbit pAb

20ng/ml, 30 min; +), HT-29 cells growing asynchronously (-) or arrested in mitosis by treatment with Thymidine (Thy; 2mM, 17 hours) followed by Nocodazole (Noc; 100ng/ml, 24 hrs; +), and NIH/3T3 cells, untreated (-) or treated with Human Platelet Derived Growth Factor- (hPDGF-AA; 50ng/ml, 30min; +) using Phospho-Stat3 (Ser727) Rabbit pAb (upper) or total Stat3 (124H6) Mouse (lower).

(upper) or total Stat3 (D3Z2G) Rabbit pAb (middle) or β -Actin (D6A8) Rabbit (lower). PC-3 cells, which do not express Stat3, was used as a negative control.

**For Research use only
IMMUNOLOGICAL SCIENCES**